

# Opportunities and Challenges for Modeling and Simulation in Development of Long-Acting Injectable Drug Products

**ACCP 2019 Annual Meeting**  
September 16, 2019

**Satish Sharan, Ph.D.**

Division of Quantitative Methods and Modeling, Office of Research and Standards,  
Office of Generic Drugs | CDER | U.S. FDA

## **Disclaimer**

*This presentation reflects the views of the author and should not be construed to represent FDA's views or policies.*

PERSONAL HEALTH

## *The Cost of Not Taking Your Medicine*



Paul Rogers

- Increased healthcare utilization (recurrence of symptoms, relapse, hospitalizations, cost)

# Long-Acting Injectable Drug Products



- Long-acting injectable (LAI) drug products are formulated to achieve extended drug release action from days to years when administered via intramuscular (IM) and subcutaneous (SC) routes.
- These products can help improve patient compliance with a better therapeutic option to treat patients who adhere poorly to frequently administered medication.

# Examples of FDA Approved Long-Acting Injectable Drug Products and Approved ANDAs



Trade Names	Ingredient	Indication	Dose Frequency	Approved Generic
ABILIFY MAINTENA KIT	ARIPIRAZOLE	Schizophrenia; bipolar I disorder	Monthly	0
ARISTADA	ARIPIRAZOLE LAUROXIL	Schizophrenia	Monthly, 6 weeks, 2 months	0
ARISTADA INITIO KIT	ARIPIRAZOLE LAUROXIL	Schizophrenia	One time	0
SUBLOCADE	BUPRENORPHINE	Opioid use disorder	Monthly	0
PROBUPHINE	BUPRENORPHINE HYDROCHLORIDE	Opioid Dependence	one time (6 months)	0
ATRIDOX	DOXYCYCLINE HYCLATE	Chronic adult periodontitis	1 week	0
BYDUREON BCISE	EXENATIDE	Improve glycemic control in type II diabetes	Weekly	0
BYDUREON...BYDUREON PEN	EXENATIDE SYNTHETIC	Improve glycemic control in type II diabetes	Weekly	0
YUTIQ	FLUOCINOLONE ACETONIDE	Chronic non-infectious uveitis affecting the posterior segment of the eye	36 months (one time)	0
ZOLADEX	GOSERELIN ACETATE	carcinoma of prostate, endometriosis, breast cancer	Monthly (4 weeks)	0
SUSTOL	GRANISETRON	Antiemetics for prevention of acute and delayed nausea and vomiting with chemotherapy	Weekly	0
LUPRON DEPOT...LUPRON DEPOT-PED	LEUPROLIDE ACETATE	Endometriosis, Fibroids, Advanced prostate cancer; children with central precocious puberty	1,3,4,6 months	0
ELIGARD	LEUPROLIDE ACETATE	Palliative treatment of advanced prostate cancer	1,3,4,6 months	0
LUPANETA PACK	LEUPROLIDE ACETATE; NORETHINDRONE ACETATE	Endometriosis	Monthly	0
DEPO-PROVERA	MEDROXYPROGESTERONE ACETATE	Prevention of Pregnancy	3 months	1
DEPO-SUBQ PROVERA 104	MEDROXYPROGESTERONE ACETATE	Prevention of pregnancy, endometriosis-associated pain	3 months	0
SINUVA	MOMETASONE FUROATE	Nasal polyps who had ethmoid surgery	3 months (one time)	0
VIVITROL	NALTREXONE	Alcohol/Opioid Dependence	Monthly (4 weeks)	0
SANDOSTATIN LAR	OCTREOTIDE ACETATE	Acromegaly, Carcinoid Tumors and Vasoactive Intestinal Peptide secreting tumors	Monthly (4 weeks)	0
ZYPREXA RELPREVV	OLANZAPINE PAMOATE	Schizophrenia	2, 4 weeks	0
INVEGA SUSTENNA	PALIPERIDONE PALMITATE	Schizophrenia, schizoaffective disorder, mood stabilizers or antidepressants	Monthly	0
INVEGA TRINZA	PALIPERIDONE PALMITATE	Schizophrenia	3 months	0
SIGNIFOR LAR KIT	PASIREOTIDE PAMOATE	Acromegaly, Cushing's Disease	4 weeks	0
PERSERIS KIT	RISPERIDONE	Schizophrenia	Monthly	0
RISPERDAL CONSTA	RISPERIDONE	Schizophrenia, Bipolar I Disorder	2 weeks	0
XYOSTED (AUTOINJECTOR)	TESTOSTERONE ENANTHATE	Testosterone replacement therapy	weekly	0
ZILRETTA	TRIAMCINOLONE ACETONIDE	Osteoarthritis pain of the knee	3 months (one time)	0
TRIPTODUR KIT	TRIPTORELIN PAMOATE	precocious puberty	24 weeks	0
TRELSTAR	TRIPTORELIN PAMOATE	Advanced prostate cancer	4/12/24 weeks	0

# Challenges in LAI Product Development and Lifecycle Management



- Long apparent half-life:
  - Longer time to reach steady state
  - Longer wash out time
- Challenging to propose relevant dosing scenarios, e.g.,
  - Impact of early, delayed or missed doses
  - Switching between formulations

# Challenges in Life Cycle Management of Long-Acting Injectable Products



- Bioequivalence (BE) Studies are challenging
- High drop out rate
- Steady state BE studies in patients (safety)
- Individual steady state attainment

# Opportunities for Modeling and Simulation in LAI Product Development




- Dosing regimen
  - Justification for dosing recommendation for missed doses
  - Impact of early, delayed or missed doses
  - Dose adjustment for special population
- Bridging results from previous studies/application
- Reducing cost, time; increasing efficiency

# Opportunities for Modeling & Simulation in Life Cycle Management



- Optimize BE study design
- Sample size
- Simulate bio-inequivalent scenarios
- Design/justify a shorter duration BE study

# Example: Paliperidone; Paliperidone Palmitate



ER suspension injectable

**NDA-22264; Invega Sustenna; every month - 2009**

**NDA-207946; Invega Trinza; every 3 month - 2015**

# INVEGA SUSTENNA Background




- Invega Sustenna is an atypical antipsychotic administered monthly for:
  - Treatment of schizophrenia in adults
  - Treatment of schizoaffective disorder in adults as monotherapy and as an adjunct to mood stabilizers or antidepressants

Indication	Initiation Dosing (deltoid)		Monthly Maintenance Dose (deltoid or gluteal)	Maximum Monthly Dose
	Day 1	Day 8		
Schizophrenia	234 mg (150 mg eq.)	156 mg (100 mg eq.)	39 (25 mg eq.) - 234 mg (150 mg eq.)	234 mg (150 mg eq.)
Schizoaffective disorder	234 mg (150 mg eq.)	156 mg (100 mg eq.)	78 mg (50 mg eq.) - 234 mg (150 mg eq.)	234 mg (150 mg eq.)


# Application of Quantitative Clinical Pharmacology in New Drug Development



## Dosing Regimen Based on Comparable Exposure



## Switch Between Risperidone LAI to PP1M



Modeling & Simulation has been effectively used to support development of LAI drug products.

# Product-Specific Guidance (PSG)



- FDA publishes PSGs to facilitate generic drug product availability and to assist the generic pharmaceutical industry with identifying the most appropriate methodology for developing drugs and generating evidence needed to support ANDA approval.
- PSGs describing the Agency's current thinking and expectations on how to develop generic drug products therapeutically equivalent (TE = PE + BE) to specific reference listed drugs.
- For two products to be considered bioequivalent, there should be no significant difference in the rate and extent of absorption of the active moiety, which are usually measured by C<sub>max</sub> (the maximum drug concentration) and AUC (the area under the concentration-time curve), respectively.

*Contains Nonbinding Recommendations*

**Draft Guidance on Paliperidone Palmitate**

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

- Active Ingredient:** Paliperidone palmitate
- Dosage Form; Route:** Extended-release suspension; intramuscular
- Recommended Studies:** One study

Type of study: (1) Parallel group, steady-state or (2) two-period, crossover steady-state  
 Strength: 156 mg/mL

Subjects: Male and nonpregnant female patients with schizophrenia or schizoaffective disorder who are already receiving a stable regimen of paliperidone palmitate extended-release suspension via the intramuscular route. Patients who are already receiving 156 mg of paliperidone injection every month would be eligible to participate in the study if continuing their established maintenance dose.

Additional comments: (1) FDA does not recommend that studies be conducted using healthy subjects or patients on a different antipsychotic treatment. (2) Both sites of injection (gluteal and deltoid) should be included in the study design for adequate site representation to support the results of the study. (3) More than three doses may be required to reach steady state. Pharmacokinetic (PK) data should be submitted to demonstrate that steady state has been reached for each individual.

**Analytes to measure (in appropriate biological fluid):** Paliperidone in plasma

Recommend steady state BE studies.

In patients due to safety concerns.

Individual steady state attainment.

# Commonly Asked Questions by ANDA Applicants



- What is the adequate number of doses to attain and confirm steady state?
- Is the proposed sample size sufficient?
- Is PK sampling scheme appropriate?

# Application of Quantitative Clinical Pharmacology for Generic Drug Development



- Models published or publicly available in NDA Clinical Pharmacology reviews.
- Key model parameters that can influence rate and extent of absorption can be identified and simulated to support alternative BE proposal.
- Virtual BE trial simulation taking formulation factors into account.

# Example: Paliperidone Palmitate 1 month Injection



ORIGINAL RESEARCH ARTICLE

Clin Pharmacokinetics 2009; 48 (9): 585-600  
© 2009 Adis Data Information

© 2009 Adis Data Information

U.S. Department of Health & Human Services

AA

## Population Pharmacokinetics of Intramuscular Paliperidone Palmitate in Patients with Schizophrenia: A Novel Once-Monthly, Long-Acting Formulation of an Atypical Antipsychotic

Mahesh N. Samtani,<sup>1</sup> An Vermeulen<sup>2</sup> and Kim Stuyckens<sup>2</sup>

- 1 Clinical Pharmacology, Advanced PK-PD Modeling & Simulation, Johnson & Johnson Pharmaceutical Research & Development, LLC, Raritan, New Jersey, USA
- 2 Clinical Pharmacology, Advanced PK-PD Modeling & Simulation, Johnson & Johnson Pharmaceutical Research & Development, a Division of Janssen Pharmaceutica NV, Beerse, Belgium

FDA U.S. FOOD & DRUG ADMINISTRATION

Follow FDA | En Español

SEARCH

Home

Dataset used Index + Validation Index + Validation

IOV Status IOV Included IOV Turned Off

Number of outliers excluded 54 54

Drug

FDA Home

Invega Sustenna

Company

Application


Approval

Persons with

- Approv
- Summz
- Officer/
- Printed
- Medica
- Chemis
- Pharmz
- Statist
- Microbi
- Clinical
- Risk As
- Propr
- Other A
- Other F
- Admini

Theta #	Parameter*	Estimate	Precision	Estimate	Precision
1	CL (L/hr)	4.95	1%	5.1	2%
2	CL - CRCL Power	0.376	3%	0.369	60%
3	V: Shift factor for Females	0.726	8%	0.778	14%
4	V (L)	391	3%	385	6%
5	V - BMI Power	0.889	1%	0.807	56%
6	KA: Shift factor for Females	0.765	7%	0.777	7%
7	KA: Shift factor for Deltoid Injection	1.23	3%	1.18	5%
8	KA x 10 <sup>3</sup> (hr <sup>-1</sup> )	0.488	2%	0.558	9%
9	KA: Age Power	0.311	14%	0.349	25%
10	KA: Injection Volume Exponent	0.359	3%	0.308	24%
11	ALAG1 or D2 (hr)	319	1%	316	1%
12	F2: Shift factor for Females	0.781	4%	0.8	4%
13	F2: Shift factor for Deltoid Injection	1.37	3%	1.37	4%
14	F2: Shift factor Deltoid 1.5 inch needle	1.54	6%	1.46	7%
15	F2	0.168	2%	0.171	4%
16	F2: BMI Power	0.642	1%	0.717	8%
17	F2: Injection Volume Exponent	0.288	1%	0.274	18%
	IV CL (CV%)	40%	2%	45%	11%
	IV V (CV%)	69%	4%	79%	13%
	IV KA (CV%)	59%	3%	70%	14%
	IV F2 (SD) <sup>†</sup>	0.064	2%	0.098	8%
	IOV CL (CV%)	26%	2%	0 FIX	N/A
	IOV V (CV%)	14%	2%	0 FIX	N/A
	IOV F2 (SD) <sup>†</sup>	0.07	2%	0 FIX	N/A
	Sigma (SD)	0.22	3%	0.27	3%
	OFV	-20238		-17934	

please Injectable Suspension



Note: Alag = Lag time for the first order absorption process; CL =clearance; Cp=plasma concentration; D2=duration of zero order input; F2=fraction of dose entering via zero order input; IM=intramuscular; KA=first order absorption rate constant; V=volume of distribution

Samtani, et.al , Clin Pharmacokinetics 2009

Clinical pharmacology review for Invega Sustenna at Drug@FDA


www.fda.gov

\* For parameter equations refer to the footer in Table 11

<sup>†</sup> IOV and IIV for F2 is computed for 100 mg eq. gluteal injection for a male subject with BMI of 26.8 kg/m<sup>2</sup>


N/A: Not applicable

# Example: Paliperidone Palmitate 1 month Injection



- Formulation dependent
- For BE assessment, key model parameters that can influence rate and extent of absorption can be modified to test alternative BE scenarios
- These parameters are generally drug dependent

# Bioequivalence Trial Simulation



- The red solid lines and shaded areas represent the median paliperidone concentrations and 90% prediction intervals.
- Blue solid lines and shaded areas represent the median paliperidone concentrations and 90% prediction intervals for hypothetical generic TEST formulation with lower bioavailability.

- Can improve efficiency of BE study:
  - Number of subjects achieving steady state
  - Sample size determination
  - How many doses will lead to attainment of steady state?
  - PK sampling time points
  - Test alternate BE scenarios
  - Probability of BE success



# Other Opportunities for Modeling and Simulation

- Propose alternate study designs for handling washout periods in the study design; reduce trial duration
- Incorporate critical quality attributes → in vitro/dissolution → in vivo dissolution/release using PBPK modeling and simulation approach

# List of M&S Grants/Contracts for LAI Products



Project title	Study duration	Grantee/Contractor	Grant/Contract No.
Development of model-informed bioequivalence evaluation strategies for long-acting injectable products	2019-2021	Uppsala University	75F40119C10018
Pharmacometric modeling and simulation for evaluation of bioequivalence for leuprolide acetate injection*	2015-2019	University of Utah	U01FD005442
Development of PBPK simulation for long-acting injectable microspheres	2015-2018	Simulations Plus Inc.	U01FD005463

Welcome to propose and submit proposals to advance regulatory science.

**GDUFA Regulatory Science:** <https://www.fda.gov/industry/generic-drug-user-fee-amendments/gdufa-regulatory-science>

# Summary

- Modeling & Simulation provides an important tool to support new and generic drug development:
  - Alternative BE approaches
  - Sample size determination
  - Effects of dose initiation; early, delayed, missed doses; switching products
  - Efficiency of BE studies, attainment of steady state; test alternative BE scenarios

# Thank You!



- Model-Informed Drug Development Pilot Program (New Drug Development)  
<https://www.fda.gov/drugs/development-resources/model-informed-drug-development-pilot-program>
- For information about Model-Informed Drug Development Pilot Program (New Drug Development) please contact [MIDD@fda.hhs.gov](mailto:MIDD@fda.hhs.gov)
- Alternative approaches to demonstrate bioequivalence: Applicants can submit their proposal through FDA's Pre-ANDA program.
- Pre-ANDA Program Information:  
<https://www.fda.gov/Drugs/ResourcesForYou/Consumers/BuyingUsingMedicineSafely/GenericDrugs/ucm578012.htm>
- For questions about submitting Pre-ANDA meeting requests for complex generic drug products online please contact [PreANDAHelp@fda.hhs.gov](mailto:PreANDAHelp@fda.hhs.gov)

# Acknowledgement



## **FDA/OMPT/CDER**

### **OGD/ORS/DQMM**

Hezhen Wang

Lanyan (Lucy) Fang

Myong-Jin Kim

Liang Zhao

QCP Team

### **OGD/ORS/DTP**

Yan Wang & Team

### **OGD/ORS-IO**

Lei Zhang

Robert Lionberger

### **OTS/OCP**

### **DQMM External Collaborators**

Uppsala University, Contract # 75F40119C10018

Simulations Plus, Inc., Grant #: U01FD005463

University of Utah Grant # U01FD005442

[www.fda.gov/GDUFARegScience](http://www.fda.gov/GDUFARegScience)



**U.S. FOOD & DRUG**  
ADMINISTRATION

# Challenge Question

Q) What are the challenges in lifecycle management of long acting injectable products?

- 1) Long durations of bioequivalence studies
- 2) Highly variable gastric emptying time
- 3) Both 1 and 2
- 4) None of the above